

# General Surgery

## ICD-10-CM 2014: Reference Mapping Card

| ICD-9-CM                                      | ICD-10-CM   |
|---|---|
| 278.01 Morbid obesity                         | E66.01 Morbid (severe) obesity due to excess calories   |
| 455.2 Internal hemorrhoids with complications | K64.8 Other hemorrhoids<br>Applicable to:<br><i>Internal hemorrhoids, without mention of degree</i><br><i>Prolapsed hemorrhoids, degree not specified</i>                               |
| 550.9 Inguinal hernia unilateral              | K40.90 Unilateral inguinal hernia, without obstruction or gangrene, not specified as recurrent<br>Applicable to:<br><i>Inguinal hernia NOS</i><br><i>Unilateral inguinal hernia NOS</i> |
| 550.92 Bilateral inguinal hernia              | K40.20 Bilateral inguinal hernia, without obstruction or gangrene, not specified as recurrent<br>Applicable to:<br><i>Bilateral inguinal hernia NOS</i>                                 |
| 553.1 Umbilical hernia                        | K42.9 Umbilical hernia, without obstruction or gangrene<br>Applicable to:<br><i>Umbilical hernia NOS</i>  |
| 553.2 Ventral hernia, unspecified             | K43.9 Ventral hernia, without obstruction or gangrene<br>Applicable to:<br><i>Epigastric hernia</i><br><i>Ventral hernia NOS</i>  |
| 553.21 Incisional hernia                      | K43.2 Incisional hernia, without obstruction or gangrene<br>Applicable to:<br><i>Incisional hernia NOS</i>  |
| 560.0 Intestinal obstruction, unspecified     | K56.60 Unspecified intestinal obstruction<br>Applicable to:<br><i>Intestinal obstruction NOS</i>  |
| 562.11 Diverticulitis of colon                | K57.32 Diverticulitis of large intestine without perforation or abscess without bleeding  |

| ICD-9-CM  | ICD-10-CM  |
|---|--|
| 565.1 Rectal/Anal fistula   | K60.3 Anal fistula<br>K60.4 Rectal fistula<br>Applicable to:<br>K60.5 Fistula of rectum to skin<br>Anorectal fistula |
| 574.1 Cholelithiasis with chronic cholecystitis                   | K80.18 Calculus of gallbladder with other cholecystitis without obstruction  |
| 574.2 Cholelithiasis without cholecystitis                        | K80.18 Calculus of gallbladder without cholecystitis without obstruction   |
| 585.5 Chronic kidney disease, stage V                             | N18.5 Chronic kidney disease, Stage 5  |
| 585.6 End state renal disease                                     | N18.6 End state renal disease<br>Applicable to:<br><i>Chronic kidney disease requiring chronic dialysis</i>          |
| 706.2 Sebaceous cyst  | L72.3 Sebaceous cyst   |
| 793.8 Abnormal mammogram, unspecified                             | R92.8 Other abnormal and inconclusive findings on diagnostic imaging of breast                                       |
| V58.31 Encounter for change or removal of surgical wound dressing | Z48.01 Encounter for change or removal of surgical wound dressing*   |

*\*Z48.01 is considered unacceptable as a principal diagnosis as it describes a circumstance which influences an individual's health status but not a current illness or injury, or the diagnosis may not be a specific manifestation but may be due to an underlying cause.*

|  |   |
|--|---|
| V67.00 Follow-up examination, following surgery, unspecified | Z09 Encounter for follow-up examination after completed treatment for conditions other than malignant neoplasm**<br>Applicable to:<br><i>Medical surveillance following completed treatment</i> |
|--|---|

**\*\* Use additional code to identify any applicable history of disease code (Z86.-, Z87.-)**  
**\*\* Z09 is considered exempt from POA reporting**

### ICD-10 Made Simple - DOCUMENT!

|                   |   |
|-------------------|---|
| <b>Acuity</b>     | Acute, chronic, intermittent  |
| <b>Severity</b>   | Mild, moderate, severe  |
| <b>Etiology</b>   | Trauma, diabetes, renal failure, exercise or infection induced  |
| <b>Location</b>   | Where is it- be specific about which joint, chest, femur, posterior thorax  |
| <b>Laterality</b> | Which side is it? Left, right, both   |
| <b>Detail</b>     | Present on admission status, associated symptom (hypoxia, loss of consciousness), additional medical diagnoses, initial versus subsequent encounter |