

Geriatric Medicine

ICD-10-CM 2014: Reference Mapping Card

Description	
*	Code is not usually sufficient justification for admission to an acute care hospital when used as a principal diagnosis.
#	Only applicable to adult patients aged 15-124 years inclusive.
%	Considered unacceptable as a principal diagnosis as it describes a circumstance which influences an individual's health status but not a current illness or injury, or the diagnosis may not be a specific manifestation but may be due to an underlying cause.
^	Considered exempt from POA reporting.

ICD-9-CM	ICD-10-CM
250.00	Diabetes mellitus without mention complication, type II or unspecified type
250.02	Diabetes mellitus without mention complication, type II or unspecified type, uncontrolled
276.1	Hyponatremia and/or hyposmolality
285.8	Anemia
332.0	Parkinson's disease
401.9	Hypertension
414.00	Coronary atherosclerosis of unspecified vessel
415.19	Other pulmonary embolism and infarction

ICD-9-CM	ICD-10-CM
424.1	Aortic valve disorder
425.4	Cardiomyopathy, other primary
427.31	Atrial fibrillation
436	Acute cerebrovascular disease
458.0	Orthostatic hypotension
482.9	Bacterial pneumonia, unspecified
571.5	Cirrhosis of liver
578.9	Gastrointestinal hemorrhage

ICD-10-CM	
I35.0	Nonrheumatic aortic (valve) stenosis
I35.1	Nonrheumatic aortic (valve) insufficiency
I35.2	Nonrheumatic aortic (valve) stenosis with insufficiency
I35.8	Other nonrheumatic aortic valve disorders
I42.5	Other restrictive cardiomyopathy
I42.8	Other cardiomyopathies
I48.91	Unspecified atrial fibrillation
I67.89	Other cerebrovascular disease
I95.1	Orthostatic hypotension
J15.9	Unspecified bacterial pneumonia
K74.0	Hepatic fibrosis
K74.69	Other cirrhosis of liver
K92.2	Gastrointestinal hemorrhage, unspecified

Geriatric Medicine

ICD-10-CM 2014: Reference Mapping Card

ICD-9-CM		ICD-10-CM	
585.6	End stage renal disease	N18.6	End stage renal disease Applicable to: Chronic kidney disease requiring chronic dialysis <i>Use additional code to identify dialysis status (Z99.2)</i>
		Z99.2	Dependence on renal dialysis % ^ Applicable to: <i>Hemodialysis status Peritoneal dialysis status Presence of arteriovenous shunt for dialysis Renal dialysis status NOS</i>
599.0	Urinary tract infection	N39.0	Urinary tract infection, site not specified <i>Use additional code (B95 – B97) to identify infectious agent</i>
682.2	Cellulitis, abscess leg	R26.2	Difficulty in walking, not elsewhere classified
724.2	Low back pain	M54.5	Low back pain Applicable to: <i>Loin pain Lumbago NOS</i>
728.87	Muscle weakness	M62.81	Muscle weakness (generalized)
781.2	Abnormality of gait	R26.0	Ataxi gait Applicable to: <i>Staggering gait</i>
		R26.1	Paralytic gait Applicable to: <i>Spastic gait</i>
		R26.89	Other abnormalities of gait and mobility
782.3	Edema	R60.0	Localized edema
		R60.1	Generalized edema
783.7	Adult failure to thrive	R62.7	Adult failure to thrive
787.2	Dysphagia, unspecified	R13.0	Aphagia Applicable to: <i>Inability to swallow</i>
790.7	Bacteremia	R78.81	Bacteremia

ICD-10 Made Simple - DOCUMENT!

Acuity	Acute, chronic, intermittent
Severity	Mild, moderate, severe
Etiology	Trauma, diabetes, renal failure, exercise or infection induced
Location	Where is it- be specific about which joint, chest, femur, posterior thorax
Laterality	Which side is it? Left, right, both
Detail	Present on admission status, associated symptom (hypoxia, loss of consciousness), additional medical diagnoses, initial versus subsequent encounter