

Gynecologic Oncology

ICD-10-CM 2014: Reference Mapping Card

Description
^ Only applicable to female patients.
Only applicable to adult patients aged 15-124 years inclusive.
% Considered exempt from POA reporting.
@ This is a specific code that can be used to specify a diagnosis.
& This is a non-specific code that cannot be used to indicate a diagnosis.
\$ Code is considered as unacceptable as a principal diagnosis as it describes a circumstance which influences an individual's health status but not a current illness or injury, or the diagnosis may not be a specific manifestation but may be due to an underlying cause.

ICD-9-CM	ICD-10-CM
180.0 Malignant neoplasm of endocervix	C53.0 Malignant neoplasm of endocervix ^
182.0 Malignant neoplasm of corpus uteri	C54.1 Malignant neoplasm of endometrium ^ C54.2 Malignant neoplasm of myometrium ^ C54.3 Malignant neoplasm of fundus uteri ^
184.0 Malignant neoplasm of vagina	C52 Malignant neoplasm of vagina ^
195.8 Malignant neoplasm site not elsewhere classified	C76.8 Malignant neoplasm of other specified ill-defined sites Applicable to: <i>Malignant neoplasm of overlapping ill-defined sites</i>
197.8 Secondary malignant neoplasm digestive organs, spleen	C78.7 Secondary malignant neoplasm of liver and intrahepatic bile duct C78.89 Secondary malignant neoplasm of other digestive organs
198.82 Secondary malignant neoplasm, genital organs	C79.82 Secondary malignant neoplasm of genital organs
233.32 Carcinoma in situ, vulva	D07.1 Carcinoma in situ of vulva ^ Applicable to: <i>Severe dysplasia of vulva Vulvar intraepithelial neoplasia III (VIN III)</i>
256.31 Premature menopause	E28.310 Symptomatic premature menopause ^ # Applicable to: <i>Symptoms such as flushing, sleeplessness, headache, lack of concentration, associated with premature menopause</i> E28.319 Asymptomatic premature menopause ^ # Applicable to: <i>Premature menopause NOS</i>
278.01 Morbid obesity	E66.01 Morbid (severe) obesity due to excess calories Use additional code to: <i>Identify body mass index (BMI), if known (Z68. -)</i>

ICD-9-CM	ICD-10-CM
617.4 Vaginal endometriosis	N80.4 Endometriosis of rectovaginal septum and vagina ^
618.5 Prolapse of vaginal vault after hysterectomy	N99.3 Prolapse of vaginal vault after hysterectomy ^
621.33 Endometrial hyperplasia with atypia	N85.02 Endometrial intraepithelial neoplasia (EIN) ^ Applicable to: <i>Endometrial hyperplasia with atypia</i>
625.8 Female genital symptoms, specified	N94.89 Other specified conditions associated with female genital organs and menstrual cycle ^
626.8 Dysfunctional uterine bleeding	N92.5 Other specified irregular menstruation ^ N93.8 Other specified abnormal uterine and vaginal bleeding ^ Applicable to: <i>Dysfunctional or functional uterine or vaginal bleeding NOS</i>
627.1 Postmenopausal bleeding	N95.0 Postmenopausal bleeding ^
627.2 Female climacteric state	N95.1 Menopausal and female climacteric states ^ Applicable to: <i>Symptoms such as flushing, sleeplessness, headache, lack of concentration, associated with natural (age-related) menopause</i>
629.1 Hydrocele canal nuck-fem	N94.89 Other specified conditions associated with female genital organs and menstrual cycle ^
654.23 Previous C-section, NOS, antepartum	O34.21 Maternal care for scar from previous cesarean delivery ^ #
729.81 Swelling of limb	M79.89 Other specified soft tissue disorders Applicable to: <i>Polyalgia</i>
788.41 Urinary frequency	R35.0 Frequency of micturition
789.07 Abdominal pain generalized	R10.84 Generalized abdominal pain
789.33 Abdominal, pelvic swelling mass, lump right lower quadrant	R19.03 Right lower quadrant abdominal swelling, mass and lump
789.34 Abdominal, pelvic swelling mass, lump left lower quadrant	R19.04 Left lower quadrant abdominal swelling, mass and lump
789.59 Other ascites	R18.8 Other ascites Applicable to: <i>Ascites NOS Peritoneal effusion (chronic)</i>

Gynecologic Oncology

ICD-10-CM 2014: Reference Mapping Card

ICD-9-CM	ICD-10-CM
793.5 Abnormal findings, genitourinary organs, radiological or other exam	R93.4 Abnormal findings on diagnostic imaging of urinary organs Applicable to: <i>Filling defect of bladder found on diagnostic imaging</i> <i>Filling defect of kidney found on diagnostic imaging</i> <i>Filling defect of ureter found on diagnostic imaging</i>
795.82 Elevated cancer antigen 125 (CA 125)	R97.1 Elevated cancer antigen 125 (CA 125) ^
998.31 Disruption of internal operation (surgical) wound	T81.32XA Disruption of internal operation (surgical) wound, not elsewhere classified, initial encounter T81.32XD Disruption of internal operation (surgical) wound, not elsewhere classified, subsequent encounter T81.32XS Disruption of internal operation (surgical) wound, not elsewhere classified, sequela encounter

ICD-9-CM	ICD-10-CM
V10.3 History of breast malignancy	Z85.3 Personal history of malignant neoplasm of breast Use additional code to identify: <i>Alcohol use and dependence (F10. - **)</i> <i>Exposure to environmental tobacco smoke (Z77.22*-)</i> <i>History of tobacco use (Z87.891*-.*)</i> <i>Occupational exposure to environmental tobacco smoke (Z57.31*-.*)</i> <i>Tobacco dependence (F17. - *-.*-*)</i> <i>Tobacco use (Z72.0*-.*-.*-*)</i>
	F10 & Z77.22 Contact with and (suspected) exposure to environmental tobacco smoke (acute) (chronic) @ Applicable to: <i>Exposure to second hand tobacco smoke (acute) (chronic)</i> <i>Passive smoking (acute) (chronic)</i>
	Z87.891 Personal history of nicotine dependence % @
	Z57.31 Occupational exposure to environmental tobacco smoke @ \$
	F17 & Z72.0 Tobacco use % @ \$ Applicable to: <i>Tobacco use NOS</i>

ICD-10 Made Simple - DOCUMENT!

Acuity	Acute, chronic, intermittent
Severity	Mild, moderate, severe
Etiology	Trauma, diabetes, renal failure, exercise or infection induced
Location	Where is it- be specific about which joint, chest, femur, posterior thorax
Laterality	Which side is it? Left, right, both
Detail	Present on admission status, associated symptoms (hypoxia, loss of consciousness), additional medical diagnoses, initial versus subsequent encounter