

Psychologist

ICD-10-CM 2014: Reference Mapping Card

ICD-9-CM		ICD-10-CM	
311	Depressive disorder	F32.9	Depressive disorder
296.2	Major depressive disorder, single episode	F32.9	Major depressive disorder, single episode
300.01	Panic disorder without agoraphobia	F40.02	Panic disorder without agoraphobia
300.02	Generalized anxiety disorder	F41.1	Generalized anxiety disorder
303.91	Alcohol dependency continuous	F10.29	Alcohol dependency continuous
300.4	Dysthymic disorder	F34.1	Dysthymic disorder
296.64	Bipolar I disorder, most current mixed, severe, specified as with psychotic behavior	F31.64	Bipolar I disorder, most current mixed, severe, specified as with psychotic behavior
V62.82	Bereavement, uncomplicated	Z63.4	Bereavement, uncomplicated
296.24	Major depressive disorder, severe	F32.2	Major depressive disorder, severe, single episode
		F33.2	Major depressive disorder, severe, recurrent episode
309.81	Posttraumatic stress disorder	F43.10	Posttraumatic stress disorder
300	Anxiety state		Anxiety state
296.44	Bipolar I disorder, recent episode manic, severe, specified as with psychotic behavior	F31.64	Bipolar I disorder, recent episode manic, severe, specified as with psychotic behavior
296.34	Major affective disorder, recurrent episode severe with psychotic behavior	F33.3	Major affective disorder, recurrent episode severe with psychotic behavior
303.93	Alcohol dependence in remission	F10.21	Alcohol dependence in remission
298.8	Reactive psychosis, NEC, NOS	F23	Reactive psychosis, NEC, NOS

ICD-10 Made Simple - DOCUMENT!

Acuity	Acute, chronic, intermittent
Severity	Mild, moderate, severe
Etiology	Trauma, diabetes, renal failure, exercise or infection induced
Location	Where is it- be specific about which joint, chest, femur, posterior thorax
Laterality	Which side is it? Left, right, both
Detail	Present on admission status, associated symptom (hypoxia, loss of consciousness), additional medical diagnoses, initial versus subsequent encounter